

# Population genetics of the Marine Otter (*Lontra felina*) along the Peruvian coast

## Field report - February 2010

Juan Valqui

Zoologisches Institut, Christian-Albrechts-Universität zu Kiel (CAU), Germany

jvalqui@zoologie.uni-kiel.de

### 1. Introduction

The historical distribution of the marine otter *Lontra felina* has been reduced due to anthropogenic factors such as hunting, habitat modification and contamination (UICN\_OSG 1998). Therefore, it is considered an endangered species by international organizations such as The World Conservation Union (IUCN), Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES), Convention on Migratory Species (CMS) and it is protected by Peruvian law.

Marine otters occur along the Pacific coast of South America from 6° S (today more probably 8°S, personal observation) to 56°S in Peru, Chile and Argentina, primarily on habitats with rocky outcroppings and caves above the water at high tide (Larivière 1998; Red List of Threatened Species Version 2009.1). Research on the population status in Peru has been limited to some basic census surveys. There is still discussion about the real population number as estimates vary from 200 to about 700 for the Peruvian coast (UICN 1976, in Castilla & Bahamondes, 1979; Apaza et al., 2004). No population genetic analyses on marine otters have been carried out, although knowledge on genetic variability and population structure is of great relevance to future conservation measures. The present study, covering the Peruvian distribution range of the marine otter, aims to provide the first molecular data for this endangered mustelid, standing as an example of non-invasive genetic research.

## 2. Results

Between December 1<sup>st</sup> 2009 and February 16<sup>th</sup> 2010 a 4-people team traveled by a 4x4 truck along the geographic distribution of *Lontra felina* at the Peruvian coast, covering a driving route of 3800 km. Between Vesique (9°12' S – 78°29'W) and Punta La Chira (16°31' S – 72°56' W), 39 locations (Figure 1) were visited collecting a total of 95 scat samples in 22 locations (Table 1). Samples of feces were collected using surgical gloves (if necessary with the help of a mechanical arm) (Figure 2) and placed into 50 ml tubes, with ethanol at 96%. The tubes were refrigerated in a cooler with ice. Date, location name, GPS position and number of samples per location were registered, as well as direct sightings and otter tracks at each location (Table 1).

22 individuals could be sighted directly, while tracks were found in 20 locations. In 6 locations, either direct sightings occurred (Figure 3) or tracks were found (Figure 4), but no scat samples could be collected (Table 1). In 11 locations no scats or tracks could be found and no otters were sighted, although local fishermen reported to see them frequently in all of them. 6 samples were collected out of 2 dead otters (skin, hair and teeth) found. One of those otters was found apparently only few days after dying and after being sampled it was donated to the Centro de Ornitología y Biodiversidad (CORBIDI) for taxidermy preparation.

From the total of 39 locations visited, eight are part of the guano administration areas, managed by the Peruvian governmental agency AGRORURAL.

Samples are momentarily held refrigerated at CORBIDI in Lima, Peru. Once the CITES authority in Peru releases the export permission, they will be sent to the Zoologisches Institut of the Christian-Albrechts-Universität zu Kiel, Germany, for DNA extraction, sequencing and further analyses.

### 3. Conclusions

- Touring 3800 km a linear distance of 1200 km at the Peruvian coast was covered.
- Except 275 km of coastline not covered in the department of Ica, each location was less than 100 km apart from the following one.
- Due to financial limitations, locations south from Punta La Chira were not visited.
- 39 locations of the geographic distribution of *L. felina* at the Peruvian coast were visited.
- 95 samples were collected at 22 locations.

### 4. Recommendations

- The results of this study should be considered in further conservation measures regarding *L. felina*.
- Similar studies should be done in the southern part of its distribution, in Chile and Argentina, to provide more solid arguments on otter population management and conservation measures.

## 5. Acknowledgments

This field work was funded by *Yaqu Pacha* and by the *Zoologisches Institut* der Christian-Albrechts-Universität zu Kiel, Germany.

The Natural Resources Agency (Dirección General Forestal y de Fauna Silvestre - DGFFS) from the Ministry of Environment of the Republic of Peru provided the authorization N°454-2009-AG-DGFFS-DGEFFS for this project. Guano administration areas were accessed with the authorization of the Peruvian governmental agency AGRORURAL.

Through the field work the following persons were of great help and are gratefully acknowledged: Armando Valdés from LEB of the Universidad Peruana Cayetano Heredia; IDEA WILD for borrowing the camera, binoculars and Walkie Talkies through Elisa Ruiz. Lucas Valqui for the mechanical arm, Ronald D´Brot and Michael Valqui for the GPS; field work team: Ronald D´Brot, Enrique Hon, Karen Verde, Celia Cáceres, Elisa Ruiz, and specially Natalia Ortiz and Jerico Solis, who also helped me with the bureaucracy. ENAPU authorities in Salaverry for the allowance to enter the piers; Cesar for the sleeping commodities in Vesique; the AGRORURAL officers: Rodolfo Beltrán and José Díaz and AGRORURAL guards: Antonio Jiménez and Zenón at Punta Colorado; Oscar Carrasco at Punta Salinas; Paulo Guerrero, Leandor Ipurre and Jaime Milla at San Juan de Marcona; Marcos Huaroto and Mauro at Punta Lomas; Pedro Agurto and Luis Domínguez at Punta Atico; Jorge Tarazona, Jose Ferrel and Lorenzo Timana at Punta La Chira; Luis and Walter Gonzales at Pachacamac Island. Karel Hartinger, Daniel Sánchez and Gianni at Ancón; Julián Durán at Pucusana; Rafael Roca at Totoritas; Olivie Wetkin from Wasipunko and Orlando at San Fernando; Thomas Valqui, Luis Alza and Wilfredo Ñáñez of CORBIDI and Ana Maria Haase for the general support on field work logistics.

## 6. References

APAZA, M.; VALQUI, J.; MANGEL, J.; ROCA, M.; ALFARO, J.; SANTILLAN, L.; PERRET, J.P.; ONTON, G.; CASTANEDA, C.; MUNEMURA, G.; TOVAR, A. (2004) Estado de Conservación de *Lontra felina* (Molina 1782) en la Costa Peruana. Reporte para la Comisión Permanente del Pacífico Sur. Lima. 23p.

CASTILLA, J. C. & BAHAMONDES, I. (1979). "Observaciones conductuales y ecológicas sobre *Lutra felina* (Molina 1782) (Carnivora: Mustelidae) en las zonas central y centro-norte de Chile." Archivos de Biología y Medicina Experimentales 12: 119-132.

LARIVIERE, S. (1998) *Lontra felina*. Mammalian Species, 575: 1-5.

Red List of Threatened Species. Version 2009.1. [www.iucnredlist.org](http://www.iucnredlist.org). Accessed 21 July 2009

UICN-OSG (1998) Simposio del Grupo especial de Nutrias de UICN. Resumen. p: 34:36.