

WORKSHOP ON THE STATUS OF SOUTH AMERICAN SEA LIONS ALONG THE DISTRIBUTION RANGE

Valparaíso, Chile, 15-17 June 2009

Organizing Committee

Doris Oliva, Enrique A. Crespo, Silvana L. Dans and Maritza Sepúlveda

INTRODUCTION AND JUSTIFICATION

The South American sea lion (*Otaria flavescens*), is one of the species of marine mammals in South America that interacts with all human activities, mainly fisheries, aquaculture and tourism. On occasions is also affected by industrial activities and oil extraction and transport. Its distribution in South America ranges from Torres in southern Brazil (29°20'S; 49°43'W; Rosas et al. 1994) to Zorritos (4°S) in Perú (Riedman 1990), passing through Cape Horn in the extreme south of the continent.

Regarding abundance and population trend there is no uniform information. One of the most studied areas is northern Patagonia in Argentina and with less quality central and southern Patagonia (Crespo, 1988; Crespo & Pedraza, 1991; Dans et al., 2004; Reyes et al., 1999; Schiavini et al., 2004). In the north the population was dramatically reduced for leather and oil between the 30's and the 50's, when a population of about 180.000 individuals fell to less of 10.000 around Península Valdés, one of the most exploited zones. Southern Chubut, Santa Cruz, Tierra del Fuego and the Falkland (Malvinas) Islands showed the same population fall, even those rookeries were less or not exploited. During the 60's a protected areas policy was developed in Chubut and the most important rookeries were protected. The population did not show any sign of recovery up to the early 90's (Crespo & Pedraza, 1991). At present this stock has increased up to 55.000 sea lions (Dans et al., 2004; Grandi et al., 2008).

In Argentina sea lions interact with gillnet fisheries in Buenos Aires Province where individual sea lions damage part of the catch biting sharks in the belly and consuming the liver but where does not get entangled (Corcuera et al., 1994; Crespo et al., 1994) The same behaviour is shown in northern Patagonia where sea lions interact with longline fisheries in Golfo San Matías (R. González, Pers.Comm.). In central Patagonia sea lions get entangled in trawling nets for hake and shrimp and an estimated figure of 200-600 individuals are taken each year (Crespo et al., 1994, 1997). While sea lions increase at a rate of 5.7% annually (Dans et al., 2004) in Uruguay decline at a rate of 4% (Paéz, 2005) showing a very local effect of mortality, on the other hand a separate population dynamics may exist for this relative restricted zone. In Uruguay as in southern Brazil sea lions are shot by fishermen as a consequence of gear damage and because there is a generalized perception that the species is a competitor for fishing resources.

As it was described for the Steller and the California sea lions, South American sea lions show different rates of increase in different rookeries. Present information about genetic units in the South-western Atlantic is partially contradictory. While some studies did not found genetic differences between rookeries from Uruguay and Península Valdés (Szapkievich et al., 1999) and between Península Valdés and the Falkland (Malvinas) Islands (Freilich, 2004), more recent studies found some degree of isolation (Túnez et al., 2006). Some individual males have been found marked in Uruguay and Buenos Aires Province breeding in Península Valdés. Feeding studies indicate that sea lions use different resources in Uruguay and Buenos Aires Province with respect to Patagonia (Koen Alonso et al., 2000; Szteren, 2006; Szteren & Páez, 2002) leading to think that presumably at least two ecological stocks could exist within the same genetic stock with gene flow based on male transport.

In the extended Chilean coast a population of around 140.000 individuals was estimated (Venegas et al. 2001, Bartheld et al. 2007, Sepúlveda et al. 2007, Oliva et al. 2008). As in Argentina the population trend is not uniform. While in northern and central-southern Chile the population is increasing, in central Chile the population is stable and in the extreme south is declining. In Chile, sea lions interact with artisanal fisheries (Oporto et al. 1991, Rodríguez 2005, Sepúlveda et al. 2007),

industrial fisheries (Hückstädt & Antezana, 2003) and salmon farms (Sepúlveda & Oliva 2005). In general only operational interactions have been considered, recording predation on fishing catches and damage to fishing gear and aquaculture installations. However, even it is known that an important part of diet includes important commercial fish species (Aguayo & Maturana 1973, George-Nascimento et al., 1985, Sielfeld et al., 1997) the information about ecological interactions is scarce (Arata & Hucke-Gaete, 2005). As a result of these conflicts and an attempt of diminishing interactions with artisanal fisheries and aquaculture activities, an Action Plan and Management of sea lions was developed in southern Chile. However up to date it has not been implemented.

In Perú, the population was subject to uncontrolled exploitation during the first half of the XX century mainly for its leather and fur (Tovar & Fuentes 1984). Exportations of fur between 1925 and 1946 raised several thousand furs per year (Majluf & Trillmich, 1981). The first surveys were carried out during the 50's (Piazza 1959 cited by Tovar & Fuentes 1984). Later, between 1968 and 1979 the first national surveys estimated a population of about 20000 individuals (Majluf & Trillmich, 1981). Since 1997 an annual record is carried out with a standardized methodology, when 144087 individuals were estimated. This figure lowered in the following year given an ENSO event in 97/98, recording 27991 individuals. In the last three national censuses carried out in 2003, 2004 y 2005 the population was estimated in 75158, 59399 and 100220 individuals, showing a decline first and an increase later in the total number.

At the light of the arguments exposed it seems important to join the sea lion specialists in the region in order to discuss the status of the sea lion population in their own countries, the most important problems they suffer and the potential solutions and courses of action. The identification of vacancy areas in knowledge will be welcome and the clarification of conservation and management policies at the national and international level.

GENERAL OBJETIVES

Assess the status of sea lion population in the distribution area (Argentina, Brazil, Chile, Falkland (Malvinas) Islands, Perú and Uruguay).

Assess regional conflicts between sea lions and human activities (fisheries, aquaculture and tourism).

Assess about administration measures applied to the species and management plans per country.

Assess about priority or vacancy areas in research for conservation and potential financial sources.

SPECIFIC OBJETIVES

Assess population trends along the distribution range.

Identify management units by means of genetic markers, ecological stocks and movements and migration.

Assess impact levels from abundance estimations, mortality rates and predation on salmon farms and fishing gears.

PARTICIPANTS INVITED

Argentina: Enrique Crespo, Silvana Dans, Florencia Grandi, Diego Rodríguez, Claudio Campagna, Gisela Giardino, Mariela Dassis

Brazil: Larissa Rosa de Oliveira

Chile: Walter Sielfeld, Doris Oliva, Maritza Sepúlveda, Layla Osmann, Anelio Aguayo, Héctor Pavéz, Roberto Schlatter, René Duran, Alejandra Urra, Constanza Weinberger, Jorge Acevedo, Francisco Ponce

Perú: Patricia Majluf

United Kingdom: David Thompson

Uruguay: Diana Szteren, Valentina Franco Trecu, Federico Riet Sapriza, Matías Feijoo

SUPPORTING AND FINANCING INSTITUTIONS

Zoo d'Amneville (France)
Yaqu-Pacha (Germany)
Dutch Zoo Conservation Fund
Subsecretaría de Pesca, Government of Chile
Latin American Society for the Study of Aquatic Mammals (SOLAMAQ)
Dirección de Investigación, Universidad de Valparaíso

REFERENCES

- Aguayo A & R Maturana. 1973. Presencia del lobo marino común *Otaria flavescens* en el litoral chileno. *Biología Pesquera* 6: 49-75.
- Aguayo A, Sepúlveda M, Palma F, Díaz H & J Yañez. 1998. Censo poblacional de lobos marinos comunes en el litoral de la V a la IX Región, 218 p. Informe final proyecto Fondo de Investigación Pesquera 96-51, Subsecretaría de Pesca, Valparaíso.
- Corcuera, J., F. Monzón, E.A. Crespo, A. Aguilar & J.A Raga. 1994. Interactions between marine mammals and coastal fisheries of Necochea and Claromecó (Buenos Aires Province, Argentina). *International Whaling Commission, (Special Issue 15):*283-290.
- Crespo, E.A. & S.N. Pedraza. 1991. Estado Actual Y Tendencia De La Poblacion De Lobos Marinos De Un Pelo (*Otaria Flavescens*) En El Litoral Patagonico. *Ecología Austral* 1: 87-95.
- Crespo, E.A., J. Corcuera & A. Lopez Cazorla. 1994. Interactions between marine mammals and fisheries in some fishing areas of the coast of Argentina. *International Whaling Commission, Special Issue 15:* 283-290.
- Crespo, E.A., S.N. Pedraza, S.L. Dans, M. Koen Alonso, L.M. Reyes, N.A. Garcia, M. Coscarella & A.C.M. Schiavini. 1997. Direct and indirect effects of the highseas fisheries on the marine mammal populations in the northern and central patagonian coast. *Journal of the Northwest Atlantic Fishery Science* 22:189-207.
- Dans, S.L., Koen Alonso, M., Crespo, E.A, Pedraza, S.N. & García, N.A. 2003. Interactions between Marine Mammals and High Seas Fisheries in Patagonia Under an Integrated Approach. Pag. 100-115 En: Gales, N., M. Hindell y R. Kirkwood (Eds) *Marine Mammals: Fisheries; Tourism and Management Issues* CSIRO Publishing
- Dans, S.L., E.A. Crespo, S.N. Pedraza & M. Koen Alonso, M. 2004. Recovery of the south american sea lion population in northern Patagonia. *Canadian J. Fisheries and Aquatic Science*. Vol. 61:1681-1690.
- Freilich, S.Y. 2004. Genetic Diversity and Population Genetic Structure in the South American Sea Lion (*Otaria flavescens*) from Argentina and the Falkland Islands. Tesis doctoral.
- George-Nascimento M, Bustamante R & C Oyarzún. 1985. Feeding ecology of the South American sea lion *Otaria flavescens*: food contents and food selectivity. *Marine Ecology Progress Series* 21: 135-143.
- Grandi M.F., Dans S.L. & Crespo E.A. 2008. Social composition and spatial distribution of colonies in an expanding population of South american sea lions. *Journal of Mammalogy* 89(5)
- Hückstädt, L.A. & T. Antezana. 2003. Behaviour of the southern sea lion (*Otaria flavescens*) and consumption of the catch during purse-seining for jack mackerel (*Trachurus symmetricus*) off central Chile. *ICES Journal of Marine Science*, 60: 1-9.
- Koen Alonso, M., EA. Crespo, SN. Pedraza, NA. García & MA. Coscarella. 2000. Food habits of the South American sea lion, *Otaria flavescens*, off Patagonia, Argentina. *Fishery Bulletin* 98: 250-263.
- Oporto J, C Mercado & L Brieva. 1991. Conflicting interactions between coastal fisheries and pinnipeds in Southern Chile, 21p. In: Report on the Benguela Ecology Programme workshop on seal-fishery biological interactions, University of Cape Town. Working paper BEP/SW91/R8.
- Oporto, J., L. Brieva, R. Navarro & Al Turner. 1999. Informe Final Proyecto FIP 97- 44. "Cuantificación poblacional de lobos marinos en la X y XI Regiones", 277 p.
- Paez, E. 2005. ¿Yo...? Otaria. ¿Y usted... III Jornadas de conservación y uso sustentable de la fauna marina.
- Reyes, L.M., Crespo E.A. & Szapkievich V. 1999. Distribution and population size of the southern sea lion (*Otaria flavescens*) in central and southern Chubut, Argentina. *Mar. Mam. Sci.* 15(2):478-493.
- Riedman, M. 1990. The pinnipeds; Seals, Sea Lions, and Walruses. Berkley: Univ. California Press.
- Rodríguez, AL. 2005. Evaluación del impacto del lobo marino común *Otaria flavescens* (Shaw, 1800) en la actividad diaria de pescadores artesanales en las caletas San Pedro de Coquimbo y Peñuelas. Tesis para obtener el grado de magíster en Ciencias del Mar. Universidad Católica del Norte – Sede Coquimbo, Chile. 98 pp.
- Rosas, F.C.W., Pinedo M.C., Marmontel M. & Haimovici M. 1994. Seasonal movements of the South American sea lion (*Otaria flavescens*, Shaw) off the Rio Grande do Sul coast, Brazil. *Mammalia* 58:51-59.
- Schiavini, A.C.M., Crespo E.A. & Szapkievich V. 2004. Status of the population of South American sea lion (*Otaria flavescens*) in Santa Cruz and Tierra del Fuego Provinces, Argentina. *Mammalian Biology (Zeit. Säuge.)* 69:1-11.

- Sepúlveda M & D. Oliva. 2005. Interactions between South American sea lions *Otaria flavescens* (Shaw) and salmon farms in southern Chile. *Aquaculture Research* 36: 1062-1068.
- Sielfeld W, C Guerra, LR Durán, E Acuña, A Aguayo, M Sepúlveda, F Palma, A Malinarich, G Cerda, A Bolvarán, R Grau, X Veloso, Y Guerra, M Vargas, N Amado, R Peredo & J Galáz. 1997. Monitoreo de la pesquería y censo del lobo marino común en el litoral de la I - IV Regiones, 105 p. Informe final Proyecto Fondo de Investigación Pesquera 95 - 28, Subsecretaría de Pesca, Valparaíso.
- Szapkievich, V.B., Cappozzo, H.L., E.A. Crespo, Bernabeu, R.O., Comas, C. & Mudry, M. 1999. Genetic relatedness in two Southern sea lion (*Otaria flavescens*) rookeries in Southwestern Atlantic. *Zeitschrift für Säugetierkunde* 64: 1-5.
- Szteren, D. 2006. Predation of *Otaria flavescens* over artisanal fisheries in Uruguay: opportunism or prey selectivity? *LAMJAM*, 5(1): 29-38.
- Szteren, D. & E. Páez. 2002. Predation by Southern sea lions (*Otaria flavescens*) on artisanal fishing catches in Uruguay. *Marine and Freshwater Research* 53: 1161-1167.
- Túnez JI, Centrón D, Cappozzo HL, & Cassini MH (2006) Geographic distribution and diversity of mitochondrial DNA haplotypes in South American sea lions (*Otaria flavescens*) and fur seals (*Arctocephalus australis*). *Mamm Biol* 72(4):193-203
- Venegas, C., J. Gibbons, A. Aguayo, W. Sielfeld, J. Acevedo, N. Amado, J. Capella, G. Guzmán & C. Valenzuela. 2001. Informe Final Proyecto FIP 2000 - 22. "Cuantificación poblacional de lobos marinos en la XII Región", 92 p.